

Suggested intervals between administration of immune globulin preparations for different indications and measles-containing vaccine and varicella vaccine*

Product/Indication	Dose, including mg immunoglobulin G (IgG)/kg body weight*	Suggested Interval before Measles or Varicella Vaccination
RSV monoclonal antibody (Synagis™)§	15 mg/kg intramuscularly (IM)	None
Tetanus (TIG)	250 units (10 mg IgG/kg) IM	3 months
Hepatitis A (IG)		
Contact prophylaxis	0.02 mL/kg (3.3 mg IgG/kg) IM	3 months
International travel	0.06 mL/kg (10 mg IgG/kg) IM	3 months
Hepatitis B IG	0.06 mL/kg (10 mg IgG/kg) IM	3 months
Rabies IG	20 IU/kg (22 mg IgG/kg) IM	4 months
Varicella IG	125 units/10kg (20-40 mg IgG/kg) IM (maximum 625 units)	5 months
Measles prophylaxis IG		
Standard (i.e., nonimmunocompromised contact)	0.25 mL/kg (40 mg IgG/kg) IM	5 months
Immunocompromised contact	0.50 mL/kg (80 mg IgG/kg) IM	6 months
Blood transfusion		
Red blood cells (RBCs), washed	10 mL/kg negligible IgG/kg intravenously (IV)	None
RBCs, adenine-saline added	10 mL/kg (10 mg IgG/kg) IV	3 months
Packed RBCs (Hct 65%)†	10 mL/kg (60 mg IgG/kg) IV	6 months
Whole blood (Hct 35-50%)†	10 mL/kg (80-100 mg IgG/kg) IV	6 months
Plasma/platelet products	10 mL/kg (160 mg IgG/kg) IV	7 months
Cytomegalovirus intravenous immune globulin (IGIV)	150 mg/kg maximum	6 months
Respiratory syncytial virus prophylaxis IGIV	750 mg/kg	9 months
IGIV		
Replacement therapy for immune deficiencies‡	300-400 mg/kg IV†	8 months
Immune thrombocytopenic purpura	400 mg/kg IV	8 months
Immune thrombocytopenic purpura	1000 mg/kg IV	10 months
Kawasaki disease	2 grams/kg IV	11 months

*This table is not intended for determining the correct indications and dosage for using immune globulin products. Unvaccinated persons might not be fully protected against measles during the entire recommended interval, and additional doses of immune globulin and/or measles vaccine might be indicated after measles exposure. Concentrations of measles antibody in an immune globulin preparation can vary by manufacturer's lot. Rates of antibody clearance after receipt of an immune globulin preparation might vary also. Recommended intervals are extrapolated from an estimated half-life of 30 days for passively acquired antibody and an observed interference with the immune response to measles vaccine for 5 months after a dose of 80 mg IgG/kg.

(Source: Mason W, Takahashi M, Schneider T. Persisting passively acquired measles antibody following gamma globulin therapy for Kawasaki disease and response to live virus vaccination [Abstract 311]. Presented at the 32nd meeting of the Interscience Conference on Antimicrobial Agents and Chemotherapy, Los Angeles, California, October, 1992.)

§Contains antibody only to respiratory syncytial virus (RSV)

†Assumes a serum IgG concentration of 16 mg/mL.

‡Measles and varicella vaccination is recommended for children with asymptomatic or mildly symptomatic human immunodeficiency virus (HIV) infection but is contraindicated for persons with severe immunosuppression from HIV or any other immunosuppressive disorder.

From ACIP "General Recommendations on Immunization" February 8, 2002